

Semi-empirical calculation of quenching factors for scintillators: new results

V.I. Tretyak

Institute for Nuclear Research, MSP 03680 Kyiv, Ukraine

Contents:

- 1. Introduction**
- 2. Outlines of the method**
- 3. Calculation of quenching factors (QF):
few old + new results**
- 4. Specific case of QF for DAMA NaI(Tl)**
- 5. Conclusions**

**Based on V.I. Tretyak, Astropart. Phys. 33 (2010) 40
+ new results**

1. Introduction

Motivations

1. Searches for dark matter:

Universe – usual matter ~4%, DM ~23%, DE ~73%;

WIMPs scatter on nuclei in a detector creating recoil ions;

Amount of light produced in scintillator by ions is lower than that produced by electrons of the same energy (experimental fact);

Thus, in scintillators calibrated with electrons or γ quanta, signals from ions will be seen at lower energies than their real values (up to ~40 times);

Evidently, knowledge of these transformation coefficients – **quenching factors (QF)** – is extremely important in searches for WIMPs: without them, you do not know where to look for the signal;

Many experimental efforts (sometimes very sophisticated) to measure QFs.

It would be very nice if one would be able to **calculate** QFs.

2. Investigation of rare ($T_{1/2}=10^{18}$ - 10^{19} yr) alpha decays with scintillators or scintillating bolometers:

Observations:

^{180}W in CdWO_4 – F.A. Danevich et al., PRC 67 (2003) 014310;

in CaWO_4 – C. Cozzini et al., PRC 70 (2004) 064606;

– Yu.G. Zdesenko et al., NIMA 538 (2005) 657;

in ZnWO_4 – P. Belli et al., NIMA 626 (2011) 31;

^{151}Eu in $\text{CaF}_2(\text{Eu})$ – P. Belli et al., NPA 789 (2007) 15;

^{209}Bi in $\text{Bi}_4\text{Ge}_3\text{O}_{12}$ – P. de Marcillac et al., Nature 422 (2003) 876;

– J.W. Beeman et al., PRL 108 (2012) 062501;

Limits:

$^{204,206,207,208}\text{Pb}$ in PbWO_4 – J.W. Beeman et al., EPJA 49 (2013) 50.


BOREXINO liquid scintillator (pseudocumene, C_9H_{12}), G. Bellini et al., PRL 107 (2011) 141302

^{210}Po

$E_\alpha = 5304 \text{ keV}$

Visible at $\sim 430 \text{ keV}$

$\text{QF}_\alpha \approx 430/5407 \approx 0.08$



**Knowledge of QF is important in all
experiments which use scintillators**

2. Outlines of the method

In calculation of QFs, we follow **Birks approach** in description of quenching of the light yield for highly ionizing particles [J.B. Birks, Proc. Phys. Soc. A 64 (1951) 874; The Theory and Practice of Scintillation Counting, Pergamon Press, Oxford, 1964]

Light yield (LY) of highly ionizing particles in scintillating material depends not only on its energy E but also on its stopping power dE/dr

Examples of stopping powers (SP) in CaWO_4 :
for *electrons* calculated with the *ESTAR* code [M.J. Berger et al., Stopping-Power and Range Tables for Electrons, Protons, and Helium Ions] and
for *different ions* calculated with the *SRIM* code [J.F. Ziegler et al., SRIM. The Stopping and Range of Ions in Matter, SRIM Co., 2008]:




Fig. 1: (a) Total SP and (b) nuclear and electronic parts of SP⁹

For particles with low SP (e^- at $E > \sim 100$ keV; S is absolute scintillation factor):

$$dL = S dE \quad \frac{dL}{dr} = S \frac{dE}{dr}$$

To account for suppression of LY for highly ionizing particles (ions) Birks proposed: (kB – Birks factor)

$$\frac{dL}{dr} = \frac{S \frac{dE}{dr}}{1 + kB \frac{dE}{dr}}$$

Approximations of light yield for electrons and ions:

$$L_e(E) = SE \quad L_i(E) = \frac{Sr}{kB}$$

but in general:

$$L(E) = \int_0^E dL = \int_0^E \frac{S dE}{1 + kB \frac{dE}{dr}}$$

Let us suppose $S_e = S_i$, and S and kB do not depend on energy

Quenching factor (often for α particles: “ α/β ratio”) is ratio of LY of ions to LY of electrons:

$$Q_i(E) = \frac{L_i(E)}{L_e(E)} = \frac{\int_0^E \frac{dE}{1 + kB(\frac{dE}{dr})_i}}{\int_0^E \frac{dE}{1 + kB(\frac{dE}{dr})_e}} \quad (1)$$

S disappeared, and **we have only 1 parameter: kB**

Sometimes instead of QF, a relative LY (ratio of ion's LY to energy, normalized to that of electron at some energy E_0) is used:

$$R_i(E) = \frac{L_i(E)/E}{L_e(E_0)/E_0} \quad (2)$$

Relation between Q and R :

(R is practically equal to Q if electron energies E and E_0 are in energy range where $L \sim E$)

$$R_i(E) = \frac{L_i(E)}{L_e(E)} \frac{L_e(E)/E}{L_e(E_0)/E_0} = Q_i(E) \frac{L_e(E)/E}{L_e(E_0)/E_0}$$

With approximations: $dL_e/dE = S$ $dL_i/dE = \frac{S}{kB} \frac{1}{(dE/dr)_i}$

we obtain the following approximation for QF:

$$Q_i(E) = \frac{L_i(E)}{L_e(E)} = \frac{L_i(E)/E}{L_e(E)/E} \simeq \frac{dL_i/dE}{dL_e/dE} \simeq \frac{1}{kB(dE/dr)_i}$$

It gives the following **important features** of QF:

- QF depends on E (in many papers constant QF was supposed);
- QF is minimal when dE/dr is maximal;
- QF increases at low E (because of decrease of dE/dr , see Fig. 1).

In the following calculations:

1. All results are obtained with Eq. (1) or Eq. (2):

$$Q_i(E) = \frac{L_i(E)}{L_e(E)} = \frac{\int_0^E \frac{dE}{1+kB(\frac{dE}{dr})_i}}{\int_0^E \frac{dE}{1+kB(\frac{dE}{dr})_e}} \quad R_i(E) = \frac{L_i(E)/E}{L_e(E_0)/E_0}$$

2. dE/dr are calculated with the **ESTAR** code for electrons and the **SRIM** code for ions (sometimes with the **ASTAR** code for α particles);

3. **Total** dE/dr is used.

Some precautions when comparing experimental and calculated QF values:

1. QF depends on kind and amount of dopant in doped scintillators (like NaI(Tl)) but also could depend on impurities and defects in “pure” scintillators (change could be tens of %);
2. QF depends on temperature (up to tens of %, see APP’2010);
3. QF depends on such a technical parameter as time Δt during which scintillation signal is collected (see Fig. 2); if signals are not collected during proper time, it is possible to obtain wrong conclusions on QF values (enhancement instead of quenching);
4. It is better to use measurements of QFs when ions’ energies are exactly known (e.g. **not** with non-monoenergetic neutron sources like Am-Be with spectrum up to ~ 11 MeV – they give mixture of QFs for different E and different ions constituting scintillator).




Fig. 2:
Data from R. Gwin, R.B. Murray,
Phys. Rev. 131 (1963) 501;
Fit by eq. (2)

**Change of Δt from $1 \mu\text{s}$ to $7 \mu\text{s}$ resulted in change of QF ($\sim 30\%$)
 and in change of kB from 1.1×10^{-3} to $2.3 \times 10^{-3} \text{ g}/(\text{MeV cm}^2)$**

**And: scintillation signal for p is faster than that for e^- ; at 662 keV
 $LY_p > LY_e$ with $\Delta t = 1 \mu\text{s}$ (thus $QF_p > 1$ – **enhancement**) while
 $LY_p < LY_e$ with $\Delta t = 7 \mu\text{s}$ (thus $QF_p < 1$ – **quenching**)**





(and sometimes Δt could be even not mentioned in a paper ...)¹⁴

In the following, we will not expect that kB is some fundamental constant of scintillating material; its value could be different in different experimental conditions (including data treatment).

However, we will suppose that if experimental conditions and data treatment are fixed, kB is the same for all particles (e^- and different ions).

Below we will check this hypothesis.

3. Calculation of QFs for different scintillators



Organic scintillators:
 C_8H_8 (polystyrene, solid)
 $C_{16}H_{18}$ (PXE, liquid)
 C_9H_{12} (pseudocumene, liquid)

For C_9H_{12} , kB values in (c) and (d) are very different – not surprise (different conditions)



In (d), kB is obtained by fitting data for protons; after this, curve for C ions is calculated

(a): M. Bongrand (SuperNEMO), AIPCP 897 (2007) 14

(b,c): H.O. Back et al. (BOREXINO), NIMA 584 (2008) 98

(d) J. Hong et al., APP 16 (2002) 333

Crystal scintillators: CdWO_4




Once more, kB values in (a) and (b) are different for the same material due to different experimental conditions.

In (b), kB value is obtained fitting data for protons; then, curve for α particles was calculated



(a): F.A. Danevich et al., PRC 67 (2003) 014310

(b): T. Fazzini et al., NIMA 410 (1998) 213

**Crystal
scintillators:
CaWO₄ – 1**



Crystal scintillators: $\text{CaWO}_4 - 2$



(a): J. Ninkovic et al.,
NIMA 564 (2006) 567.
(b): G. Angloher et al.,
APP 23 (2005) 325.


LY/E of e^- (for $E=6 \text{ keV}$) to LY/E of different ions (for $E=18 \text{ keV}$)

In (a), kB was obtained normalizing th. and exp. R' values for protons; after this, R' values for all other ions were calculated. Range of A, Z values: $A=1, Z=1$ for p, and $A=197, Z=79$ for Au.

In (a) – room T , in (b,c) $T=7 \text{ mK}$ – so, different kB .

In (b), kB was obtained to reproduce R for 2.3 MeV α particle ($A=4, Z=2$). In (c), calculated curve with this kB is in agreement with point for 104 keV Pb ions ($A=206, Z=82$).

Crystal scintillators: CsI(Tl)



QF values for Cs and I ions are practically the same.

kB is the same for measurements by different groups (this could be just coincidence).

kB for α particles is different in (d) but data were measured in different conditions.

(a): S. Pecourt et al., APP 11 (1999) 457



(b): H. Park et al., NIMA 491 (2002) 460

(c): M.Z. Wang et al., PLB 536 (2002) 203

(d): T.Y. Kim et al., NIMA 500 (2003) 337

Y.F. Zhu et al., NIMA 557 (2006) 490

**Crystal
scintillators:
CsI(Na)**




In (a), QF for Cs (or I) ions are described.

In (b), QF for α particles with $kB=5.5e-3$ is predicted.

Data: H. Park et al., NIMA 491 (2002) 460

Crystal scintillators: NaI(Tl)




(a): D.R. Tovey et al.,
PLB 433 (1998) 150

(b): H. Chagani et al.,
JINST 3 (2008) P06003





In (a), kB was obtained by describing data for Na ions and then used to calculate curve for I ions (nice agreement).

kB in (a) and in (b) are different due to different experimental conditions.



**Crystal
scintillators:
 CeF_3**

Liquid noble gases: LXe – 1




Some experimental data are well described in the current approach; different kB values are due to different conditions.

(a): R. Bernabei et al., PLB 436 (1998) 379
 (b): V. Chepel et al., APP 26 (2006) 58

(c): R. Bernabei et al., EPJ Cdir 11 (2001) 1
 (d): M. Tanaka et al., NIMA 457 (2001) 454

**Liquid
noble gases:
LXe – 2**



However, for Xe ions in LXe exist also other experimental data which **are not described** in the current approach.

Data (and summary): G. Plante et al., PRC 84 (2011) 045805


Few calculations fulfilled
after the article: V.I. Tretyak, Astropart. Phys. 33 (2010) 40
are given below

Ar ions in liquid Ar - D. Gastler et al., PRC 85 (2012) 065811:


Description of the data just by constant after 20 keV



Page 7: “An observed upturn in the scintillation efficiency below 20 keVr is currently unexplained.”




Description here in accordance with Eq. (2):




Ar ions in liquid Ar – C. Regenfus et al., arXiv:1203.0849 = JPCS 375 (2012) 012019

Experimental data with
2 models:

1. J. Lindhard et al.,
Mat. Fys. Medd. Dan.
Vid. Selsk. 33/14 (1963) 1
2. D. Mei et al., APP 30 (2008) 12





Description here in
accordance with Eq. (2):





Alpha particles in CdWO_4 - C. Arnaboldi et al., APP 34 (2010) 143

Data presented in (a) were used to obtain the kB value.

LY curve for continuous spectrum of alpha particles was calculated with this kB .



Different ions in CdWO_4 - P.G. Bizzeti et al., NIMA 696 (2012) 144




Value of kB was obtained by fitting data for protons.

Quenching curves for other ions (α , Li, C, O, Ti) were calculated with this kB , in quite good agreement with the experimental data.

Alpha particles in plastic - X. Sarazin, Memoire d'habilitation, 2012

BiPo-3: C_8H_8
(polystyrene)

Red line is calculation of
 $1/QF$




Protons in pseudocumene (C_9H_{12}) – G. Bellini et al., PRC 81 (2010) 034317

Data for alpha particles are drawn as in APP'2010



Data for protons appeared after APP'2010.

Curve QF_p for protons is calculated with the same kB value as for alpha's.

Agreement is excellent.



4. Specific case of QF for DAMA NaI(Tl)



In (c), kB is obtained by fitting data for α particles (internal contamination) in LIBRA experiment (R. Bernabei et al., NIMA 592 (2008) 297) – in the same conditions as DM data. Predicted with this kB QFs for Na and I ions in (d) are much higher ($QF_{\text{Na}}=0.64$ at 5 keV) than those usually measured and used in NaI(Tl) dark matter experiments.

C. Arina, J. Hamann, Y.Y.Y. Wong, JCAP 09 (2011) 022:

(see also JPCS 375 (2012) 012009 and 1210.4011)

Limitation: $QF_{Na} \leq 0.6$

Combined fit of the DAMA and CoGeNT, QF_{Na} is free parameter.

“If we demand compatibility between these experiments, then the inference process naturally concludes that a high value for the sodium quenching factor for DAMA is preferred.”





Figure 13. Same as figure 11, but for an extended prior range for the DAMA sodium quenching factor q_{Na} (up to $q_{Na} = 0.6$). 6

Consequence of bigger QF_{Na} and QF_I :
shift of WIMPs mass to lower values ~ 10 GeV.

P. Belli et al., PRD 84 (2011) 055014, Fig. 1 (for some set of parameters):




5. Conclusions

- (1) Old Birks formula still gives nice description of QF for ions in many cases – if *total* SP for electrons and ions are used, and SP are calculated with the **ESTAR** and **SRIM** codes which are: (a) publicly available, (b) are ones of the best codes in this field.
- (2) There is only one free parameter in the approach – the Birks kB factor. It is **not considered as some fundamental constant** for a given scintillating material but as a variable which **depends on conditions of measurements and data treatment.**
- (3) There are experimental data which confirm the hypothesis that, **once conditions of measurements and data treatment are fixed, the kB value is the same for different ions.** Thus, if kB was determined by fitting data for particles of one kind (e.g. α particles of few MeV from internal contamination), it can be used to calculate QFs for particles of another kind and for another energies of interest (e.g. low energy recoils after scattering of DM particles).

- (4) Quenching factors for ions calculated in the present approach in general **increase at low energies**, and this encourages experimental searches for DM particles.
- (5) For the DAMA/LIBRA experiment, it was shown that, based on measured in DAMA/LIBRA QFs for α particles, **QFs for Na and I ions should be ~2 higher** than those typically used in NaI(Tl) DM experiments. It shifts the “evidence spot” of the DAMA/LIBRA observations to WIMPs’ lower masses (~ 10 GeV) relaxing contradictions with other experiments which give only limits for WIMPs cross-sections.

Thank you for attention!

**5. Possible relation between QF for low energy electrons
(and γ quanta) and QF for ions**




SP for electrons is higher at low energies: so, **quenching (non-proportionality) for low energy electrons should exist?**

Answer: yes, it exists, and this fact is known many years.


For some scintillators, it is possible to describe R_e on the basis of the Birks formula:

$$R_e(E) = \frac{L_e(E)/E}{L_e(E_0)/E_0}$$


$$L_e(E) = \int_0^E dL_e = \int_0^E \frac{SdE}{1 + kB(\frac{dE}{dr})_e}$$



**CaWO₄ – R.F. Lang et al.,
arXiv: 0910.4414**





**BOREXINO liquid
scintillator – H.O. Back
et al., PLB B 525 (2002) 29**



**Double Chooz liquid
scintillator – C. Aberle
et al., JINST 06 (2011)
P11006**

However, life is more rich: there are also scintillators for which R_e cannot be described with the Birks law. Moreover, with the Birks formula it is possible to obtain only quenching (at most, constant at $kB=0$), however, there are scintillators where instead enhancement of R_e is observed at low energies.



CdWO₄ – P.G. Bizzeti et al., NIMA A 696 (2012) 144

Summary for different scintillators – S.A. Payne et al., IEEE TNS 58 (2011) 3392

So, in my current understanding, if even non-proportional behaviour of relative LY for electrons can be described on the basis of the Birks law, value of kB factor is not necessary equal to kB for description of QF for ions.

Some data on quenching of ionization signal in Ge detectors

1. α particles in Ge: S. Fiorucci et al., *Astropart. Phys.* 28 (2007) 143:
 $Q=0.30\pm 0.02$ at $E_\alpha=5.33$ MeV and $T=17$ mK

2. Ge ions in Ge: A. Benoit et al., *NIMA* 577 (2007) 558




Fig. 2. Experimental results of the direct measurement of the ionization quenching for germanium recoils in germanium, from Refs. [5,7–11]. The line represents Eq. (1), with parameter values as of Eqs. (2)–(4).

3. α particles in Ge (at $T=77$ K):
Ph. Hubert et al., *NIMA* 252 (1986) 87
also Ph. Hubert, private comm. (2007)
also G. Heusser, private comm. (2007)
also our Ge measurements in LNGS
 $Q\cong 1$ at $E_\alpha=5.33$ MeV

Puzzle ? (different electric fields ?)

Some data on quenching of ionization signal in Si detectors

Si ions in Si:

1. G. Gerbier et al., PRD 42 (1990) 3211




FIG. 4. Ratio between the observed energy [equivalent electron energy (EEE)] and the calculated recoil energy as a function of the silicon recoil energy. Circles are data points from the present experiment, squares are data points from Sattler's experiment (Ref. 8). The curve represents the result of the calculation of Lindhard *et al.* (Ref. 6).

2. A.R. Sattler, Phys. Rev. A
138 (1965) 1815




FIG. 5. Pulse height produced by a Si recoil atom relative to that of an electron of the same energy in Si as a function of Si recoil energy. The incident monoenergetic neutron energy necessary to produce the denoted recoil energy in a backscattering event is shown in parenthesis. Solid line denotes predictions of Lindhard *et al.*, in variables $\bar{\eta}/E$.

Theoretical attempts:

- 1. J.B. Birks, Proc. Phys. Soc. A 64 (1951) 874;
The Theory and Practice of Scintillation Counting, Pergamon Press,
Oxford, 1964.**
- 2. R.B. Murray, A. Meyer, Phys. Rev. 122 (1961) 815.**
- 3. J. Lindhard et al., Mat. Fys. Medd. K. Dan. Vidensk. Selsk. 33 (1963) 1
(practical receipt is reproduced in: D.-M. Mei et al., Astropart. Phys. 30
(2008) 12).**
- 4. A. Hitachi, Astropart. Phys. 24 (2005) 247;
J. Phys.: Conf. Ser. 65 (2007) 012013.**